

Fig 6 Experimental substrate concentration profile model with 150 g/l initial substrate concentration




Fig 7 Experimental gluconic acid concentration profile and model with 150 g/l initial substrate concentration

V. CONCLUSION

Batch fermentation of glucose to produce gluconic acid has been accomplished using three different substrate concentrations with high initial *Aspergillus niger* mycelium concentration. Kinetic parameters for cell growth, substrate consumption, and product formation have been determined for gluconic acid batch fermentation by *Aspergillus niger* FNCC 6098.

ACKNOWLEDGMENT

This work was supported by the Indonesian Directorate General of Higher Education (DGHE), Ministry of National Education (MONE) through Young Scientist Research Award

(Penelitian Dosen Muda No. 246/SP2H/PP/DP2M/III/2008).

REFERENCES

- [1] J.Z. Liu, L.P. Weng, Q.L. Zhang, H. Xu, L. N. Ji, "A mathematical model for gluconic acid fermentation by *Aspergillus niger*", *Biochemical Engineering Journal*, vol. 14, pp. 15-64, 2003.
- [2] S. Velizarov, V. Beschkov, "Biotransformation of glucose to free gluconic acid by *Gluconobacter oxydans*: substrate and product inhibition situations", *Process Biochemistry*, vol. 33, no. 5, pp.527-534, 1998.
- [3] H. Znad, J. Markos, V. Bales, "Production of gluconic acid from glucose by *Aspergillus niger*: growth and non-growth conditions", *Process Biochemistry*, vol. 39, pp. 1341-1345, 2004.
- [4] M.L. Shuler, F. Kargi, *Bioprocess Engineering Basic Concept*. New Jersey: Prentice-Hall Inc., 1992, pp.176.
- [5] T. Takamatsu, S. Shioya, T. Furuya, "Mathematical model of gluconic acid fermentation by *Aspergillus niger*", *J. Chem. Technol. Biotechnol.*, vol. 31, pp. 697-704, 1981.
- [6] H. Znad, M. Blazej, V. Bales, J. Markos, "A kinetic model for gluconic acid production by *Aspergillus niger*", *Chem. Pap.*, vol. 58, no. 1, pp. 23-28, 2004.